

Locally Euclidean metrics on $\mathbf{R}^3$

Young Deuk Kim
 School of Mathematical Sciences
 Seoul National University
 Seoul, 151-747, Korea
 (ydkimus@yahoo.com)

February 2, 2008

Abstract

For all $0 < t \leq 1$, we define a locally Euclidean metric ρ_t on $\mathbf{R}^3$. These metrics are invariant under Euclidean isometries and, if t increases to 1, converge to the Euclidean metric d_E . This research is motivated by expanding universe.

key words. locally Euclidean metric

PACS number(s). 98.80.Jk

Mathematics Subject Classifications (2000). 85A40, 57M50

1 The metric ρ_t

Let d_E denote the Euclidean metric on $\mathbf{R}^3$. As in [5], a nonnegative function $d : \mathbf{R}^3 \times \mathbf{R}^3 \rightarrow \mathbf{R}$ is called a *locally Euclidean metric* if

- (i) $d(P, Q) = 0$ if and only if $P = Q$
- (ii) $d(P, Q) = d(Q, P)$ for all $P, Q \in \mathbf{R}^3$
- (iii) $d(P, Q) + d(Q, R) \geq d(P, R)$ for all $P, Q, R \in \mathbf{R}^3$
- (iv) For all $P \in \mathbf{R}^3$, there exists $r > 0$ such that

$$d(Q, R) = d_E(Q, R) \quad \text{for all } Q, R \in B_r(P) = \{S \in \mathbf{R}^3 \mid d(P, S) < r\}.$$

Let $S^2 \subset \mathbf{R}^3$ be the unit 2-sphere with center $O = (0, 0, 0)$ and radius 1. A locally Euclidean metric on S^2 is defined in the same way as on $\mathbf{R}^3$. In the following theorem of the author, $-P$ is the antipodal point of $P \in S^2$ and $0 \leq \alpha < \pi/4$.

Theorem 1 ([5]) Suppose that $0 < t \leq 1$ and $\alpha = \sin^{-1} \left(\frac{\sqrt{2-t^2}-t}{2} \right)$. For all $P, Q \in S^2$ we define

$$d_t(P, Q) = \begin{cases} d_E(P, Q) & \text{if } \angle POQ \leq \pi - 2\alpha \\ 2t + d_E(-P, Q) & \text{if } \angle POQ > \pi - 2\alpha. \end{cases}$$

Then d_t is a locally Euclidean metric on S^2 which is invariant under any Euclidean isometry and

$$td_E(P, Q) \leq d_t(P, Q) \leq d_E(P, Q). \quad (1)$$

Notice that

$$\text{if } d_E(P, Q) = d_E(R, S) \text{ then } d_t(P, Q) = d_t(R, S) \quad (2)$$

and

$$\text{if } d_t(P, Q) < 2t \text{ then } d_t(P, Q) = d_E(P, Q). \quad (3)$$

We will make use of the metric d_t to define a locally Euclidean metric ρ_t on $\mathbf{R}^3$. Suppose that $P, Q \in \mathbf{R}^3$ and $n \in \mathbf{N}$. Let

$$\begin{aligned} \Gamma_{P,Q}^n &= \{ (X_0, X_1, \dots, X_n) \mid X_0 = P, X_n = Q \\ &\text{and } X_i \in \mathbf{R}^3, 0 \leq d_E(X_{i-1}, X_i) \leq 2 \text{ for all } 1 \leq i \leq n \} \end{aligned}$$

and

$$\Gamma_{P,Q} = \bigcup_{n \in \mathbf{N}} \Gamma_{P,Q}^n.$$

Notice that $\Gamma_{P,Q} \neq \emptyset$ for all $P, Q \in \mathbf{R}^3$ and if $(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}$ then $(X_n, X_{n-1}, \dots, X_0) \in \Gamma_{Q,P}$.

Suppose that $(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}$ and $1 \leq i \leq n$. There exists an isometric embedding $\epsilon_i : (S^2, d_E) \rightarrow (\mathbf{R}^3, d_E)$ such that $X_{i-1}, X_i \in \epsilon_i(S^2)$. Notice that

$$d_t(\epsilon_i^{-1}(X_{i-1}), \epsilon_i^{-1}(X_i)) \quad (4)$$

does not depend on the choice of ϵ_i because d_t is invariant under Euclidean isometries. Therefore for the sake of simplicity, we will simply write

$$d_t(X_{i-1}, X_i)$$

to denote eq. (4). In the following definition, the infimum is taken for all elements of $\Gamma_{P,Q}$.

Definition 2 For $0 < t \leq 1$ and $P, Q \in \mathbf{R}^3$, we define

$$\rho_t(P, Q) = \inf_{(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}} \sum_{i=1}^n d_t(X_{i-1}, X_i).$$

We will prove the following lemma and theorem in the next section.

Lemma 3 $td_E(P, Q) \leq \rho_t(P, Q) \leq d_E(P, Q)$ for all $P, Q \in \mathbf{R}^3$.

Theorem 4 ρ_t defines a locally Euclidean metric on $\mathbf{R}^3$.

Suppose that $\phi : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ is an Euclidean isometry. Notice that $(X_0, X_1, \dots, X_n) \in \Gamma_{P, Q}$ if and only if $(\phi(X_0), \phi(X_1), \dots, \phi(X_n)) \in \Gamma_{\phi(P), \phi(Q)}$. Therefore, from eq. (2) we can show that

$$\rho_t(P, Q) = \rho_t(\phi(P), \phi(Q)).$$

By Lemma 3 if t increases to 1, then

$$\rho_t(P, Q) \text{ converges to } d_E(P, Q) \quad \text{for all } P, Q \in \mathbf{R}^3.$$

The following corollary which will be proven in the next section implies that ρ_t and d_E are same locally.

Corollary 5 If $d_E(P, Q) < 2t$ then $\rho_t(P, Q) = d_E(P, Q)$ for all $P, Q \in \mathbf{R}^3$.

The construction of ρ_t is motivated by *expanding universe*. See [1], [2] and [3] as references on this topic. Notice that $(\mathbf{R}^3, \rho_t)$ is homogeneous, isotropic and homeomorphic to the Euclidean space $(\mathbf{R}^3, d_E)$ for all $0 < t \leq 1$. From the proof of Theorem 4 in the next section, notice also that if t increases, so is the Euclidean region $B_t(P)$ for all $P \in \mathbf{R}^3$.

The Einstein's field equations are expressed in terms of Riemannian metrics (see [6], [7] and [8] for details). If a Riemannian metric space is locally Euclidean, then it is the Euclidean space itself. Therefore the construction of this paper is impossible with Riemannian metrics. In a preprint of the author([4]), there exists a similar construction of pseudo metrics on the closed ball B^3 .

2 Proof of the lemma, theorem and corollary

Proof of Lemma 3. Suppose that $P, Q \in \mathbf{R}^3$. Choose $m \in \{0\} \cup \mathbf{N}$ such that $m \leq d_E(P, Q) < m + 1$. Find $(X_0, X_1, \dots, X_{m+1}) \in \Gamma_{P, Q}$ such that

$$X_i \in \overline{PQ} \quad \text{and} \quad d_E(X_{i-1}, X_i) = \frac{1}{m+1} d_E(P, Q) \quad \text{for all } 1 \leq i \leq m+1,$$

where $\overline{PQ}$ is the Euclidean segment from P to Q . From eq (1), we have

$$\begin{aligned} \rho_t(P, Q) &\leq \sum_{i=1}^{m+1} d_t(X_{i-1}, X_i) \leq \sum_{i=1}^{m+1} d_E(X_{i-1}, X_i) \\ &= \sum_{i=1}^{m+1} \frac{1}{m+1} d_E(P, Q) = d_E(P, Q). \end{aligned}$$

For all $(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}$, from eq. (1) we have

$$\sum_{i=1}^n d_t(X_{i-1}, X_i) \geq \sum_{i=1}^n t d_E(X_{i-1}, X_i) = t \sum_{i=1}^n d_E(X_{i-1}, X_i) \geq t d_E(P, Q).$$

Thus $\rho_t(P, Q) \geq t d_E(P, Q)$. ■

Proof of Theorem 4. Since d_t is nonnegative, so is ρ_t . By Lemma 3, it is trivial to show that $\rho_t(P, Q) = 0$ if and only if $P = Q$. Since $(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}$ if and only if $(X_n, X_{n-1}, \dots, X_0) \in \Gamma_{Q,P}$, from eq. (2) we have

$$\rho_t(P, Q) = \rho_t(Q, P) \quad \text{for all } P, Q \in \mathbf{R}^3.$$

For all $(X_0, X_1, \dots, X_n) \in \Gamma_{P,Q}$ and $(Y_0, Y_1, \dots, Y_m) \in \Gamma_{Q,R}$, we have

$$(X_0, X_1, \dots, X_n, Y_0, Y_1, \dots, Y_m) \in \Gamma_{P,R}.$$

Therefore

$$\rho_t(P, R) \leq \sum_{i=1}^n d_t(X_{i-1}, X_i) + \sum_{i=1}^m d_t(Y_{i-1}, Y_i)$$

and hence

$$\rho_t(P, R) \leq \rho_t(P, Q) + \rho_t(Q, R) \quad \text{for all } P, Q, R \in \mathbf{R}^3. \quad (5)$$

Suppose that $P \in \mathbf{R}^3$ and $Q, R \in B_t(P) = \{S \in \mathbf{R}^3 \mid \rho_t(P, S) < t\}$. From eq. (5), we have $\rho_t(Q, R) \leq \rho_t(Q, P) + \rho_t(P, R) < 2t$. Therefore there exists $(X_0, X_1, \dots, X_n) \in \Gamma_{Q,R}$ such that

$$\sum_{i=1}^n d_t(X_{i-1}, X_i) < 2t.$$

From eq. (3), we have $d_t(X_{i-1}, X_i) = d_E(X_{i-1}, X_i)$ for all $1 \leq i \leq n$. Therefore

$$\begin{aligned} \rho_t(Q, R) &= \inf_{\Gamma_{Q,R}} \sum_{i=1}^n d_t(X_{i-1}, X_i) = \inf_{\Gamma_{Q,R}(t)} \sum_{i=1}^n d_t(X_{i-1}, X_i) \\ &= \inf_{\Gamma_{Q,R}(t)} \sum_{i=1}^n d_E(X_{i-1}, X_i) \geq d_E(Q, R), \end{aligned}$$

where

$$\Gamma_{Q,R}(t) = \{ (X_0, X_1, \dots, X_n) \in \Gamma_{P,Q} \mid \sum_{i=1}^n d_t(X_{i-1}, X_i) < 2t \}.$$

Thus from Lemma 3, we have $\rho_t(Q, R) = d_E(Q, R)$. ■

Proof of Corollary 5. Suppose that $d_E(P, Q) < 2t$. From Lemma 3, we have

$$\rho_t(P, Q) < 2t. \quad (6)$$

In the last paragraph of the proof of Theorem 4, we showed that eq. (6) implies $\rho_t(P, Q) = d_E(P, Q)$. ■

References

- [1] J.D. Barrow, *Relativistic cosmology*, The early universe (eds. W. G. Unruh and G. W. Semenoff), 125-201, Reidel, Dordrecht, 1988.
- [2] G.F.R. Ellis, *The expanding universe: A history of cosmology from 1917 to 1960*, Einstein and the history of general relativity (eds. D. Howard and J. Stachel), 367-431, Birkhäuser, Boston, 1989.
- [3] M. Kafatos, *The problem of observation in cosmology and the big bang*, Gravitation and cosmology: From the Hubble radius to the Plank scale (eds. R.L. Amoroso, G. Hunter, M. Kafatos and J.P. Vigiér), 65-80, Kluwer Academic Publishers, Dordrecht, 2002.
- [4] Y.D. Kim, *A family of pseudo metrics on B^3 and its application*, preprint, arXiv:math.GN/0201077.
- [5] Y.D. Kim, *Locally Euclidean metrics on S^2 in which some open balls are not connected*, Rocky Mountain Journal of Mathematics **36**(2006), no.6, 1927-1935.
- [6] J. Norton, *How Einstein found his field equations, 1912-1915*, Einstein and the history of general relativity (eds. D. Howard and J. Stachel), 101-159, Birkhäuser, Boston, 1989.
- [7] H.C. Ohanian, *Gravitation and spacetime*, W. W. Norton & Company, New York, 1976.
- [8] C. Ray, *The cosmological constant: Einstein's greatest mistake?*, Stud. Hist. Philos. Sci. **21**(1990), no.4, 589-604.